

COURSE CODE: PRM 201
COURSE TITLE: INTRODUCTION TO FORAGE SCIENCE
NUMBER OF UNITS: 2 units
COURSE DURATION:

COURSE DETAILS:

Course Coordinator: PROF (MRS) A. O. JOLAOSHO *B. Sc., M. Sc., Ph.D.*
Email: ajolaosho@yahoo.com ; ajolaosho@gmail.com
Office Location: Pasture & Range Management Office
Other Lecturers:

COURSE CONTENT:

- IMPORTANCE OF PASTURE IN ANIMAL AND CROP PRODUCTION SYSTEMS, LEGUMES, GRASSES, BROWSE PLANTS
- CLASSIFICATION OF PASTURE GRASSES AND LEGUMES.
- ESSENTIAL SOIL NUTRIENTS FOR PASTURE PRODUCTION
- TERMINOLOGIES IN PASTURE PRODUCTION.
- PROSPECTS IN PASTURE PRODUCTION – TURF MAKING, CONSERVATION, ENGINEERING, BREEDING, PUBLIC AND PRIVATE SECTORS, ETC

READING LIST:

1. WHITEMAN, P.C. 1980: TROPICAL PASTURE SCIENCE. PUBLISHED BY OXFORD UNIVERSITY PRESS.
2. HOGSON, J. AND ILLIUS, A. W. 1996: THE ECOLOGY AND MANAGEMENT OF GRAZING SYSTEMS. CAB INTERNATIONAL WALLINGFORD, USA.

3. MICHEAL, B. JONES AND ALEC LAZENBY, 1988: THE GRASS CROP. THE PHYSIOLOGICAL BASIS OF PRODUCTION CHAPMAN AND HALL LTD USA.
4. PEARSON, C. J. AND ISN, R. L. 1987: AGRONOMY OF GRASSLAND SYSTEMS. CAMBRIDGE UNIVERSITY PRESS NY.
5. CROWDER, L. V. AND CHHEDA, H.R. 1982. TROPICAL GRASSLAND HUSBANDARY. TROPICAL AGRICULTURE SERIES. LONGMAN, LONDON.
6. HUMPHERYS, L. R. 1991: TROPICAL PASTURE UTILIZATION. PUBLISHED BY THE PRESS SYNDICATE OF THE UNIVERSITY OF CAMBRIDGE NY.
7. BOGDAN, A. V. 1977: TROPICAL PASTURE & FODDER PLANTS (GRASSES & LEGUMES). TROPICAL AGRICULTURE SERIES. LONGMAN LONDON

LECTURE NOTES

BOTANY AND SYSTEMATICS OF GRASSES AND LEGUMES

Grasses belongs to family Gramineae (Poaceae). They are made up of two main parts: shoot or tillers (aerial parts) and roots (subterranean parts) with the reproductive parts included at flowering. They are MONOCOTYLEDONOUS i.e. possess one seed leaf. They are HERBACEOUS (non-woody), divergent in size, shape and growth habit, can be annual or perennial in life form.

A. VEGETATIVE ORGANS

1. AERIAL PARTS

The shoots consists of the stem (culm, haulm) and the leaves. The **stems** are cylindrical or rounded and jointed i.e made up of **nodes** separated by **internodes**. Internodes can be hollow (e.g. *Brachiaria mutica*) filled with white pith (e.g. *Zea mays*, *Sorghum vulgare* and *Hyparrhenia spp.*) or solid e.g. *Axonopus scoparius*). stems can be glabrous or pubescent. Shoots develop from buds found at the nodes and produce side branches.

The basal portion of tufted grasses is called CROWN. Stolons are creeping stems that grow above the surface of the ground and develop roots and shoots at the nodes. Examples of grasses with stolons are *Pennisetum clandestinum*, *Cynodon nlemfuensis* and *Digitaria pentzii*.

The leaves consists of the sheath, ligule and the leaf blade. leaf blades are borne on sheaths. Leaves are situated on stem in opposite rows alternatively. The leaf sheath have free edges and chlorophyll. Leaves could be smooth or rough


The Leaf blade can be setaceous (wiry and bristle) or filiform (thread-like). The midrib is usually prominent with faint lateral veins.

LIGULE is an appendage found at the junction of the leaf blade and the sheath, usually closely ad pressed to the culm. It may be a membrane, a fringe of hairs or a hardened ring and varies in size, shape and texture, can be used for identification.

AURICLES are earlike outgrowths at the leaf base of some spp. They can be prominent and encircle the stem, minute and inconspicuous or absent, no chlorophyll, no function, but use in taxonomic identification.

COLLAR: marks the junction of the outer surface (upper region) of the sheath and leaf blade, usually discoloured, leaves breaks at the collar.

PROPHYLLUM: is a two-keeled organ (a reduced leaf) covering the bud in the axil of the sheath.



Auricles



Present



Blunt



Absent

Ligule



Absent



Membranous

Sheath


- Fused or overlapping



Fused



Overlapping



2. SUBTERRANEAN PARTS

Roots appear first from the germinating seed and are known as the PRIMARY ROOTS. Primary roots can branch and provide seedlings with water and mineral nutrients in the first stage of growth. They are then replaced with secondary roots which can be very numerous, which helps retain soil particles to prevent erosion. SECONDARY OR ADVENTITIOUS ROOTS are those that developed from the nodes of tillers or creeping stems.

B. REPRODUCTIVE ORGANS

The Floral Organs are MODIFIED SHOOTS, consisting of STAMENS and PISTILS. The floral organs are called inflorescence.

INFLORESCENCE: The flowers or inflorescences may be terminal, or axillary. The basic unit of the inflorescence is the SPIKELET, which consists of flowers usually occurring in groups or clusters.


SPIKELET: A typical spikelet consists of an axis (rachilla), two glumes and one to many florets. The perfectly developed floret has a lemma and a palea (lower and upper bracts, respectively) which enclose the flower. The structure of the lamina is such that it provides protection for the seeds and perhaps means of dispersal. The palea is shorter than the lemma and thinner.

Inflorescence types are classified as:

1. SPIKE: the spikelets are sessile (without stalks) or nearly so, on an UNBRANCHED axis (rachis) eg. *Lolium*, *Triticum*, *Secale*, *Hordeum*, *Agropyron* or be ONE-SIDED eg. *Ctenium elegans* or DIGITATE (finger like) as in *Chloris* and *Cynodon* Spp. or RACEMOSE on a central axis as in *Dactyloctenium* and *Leptochloa* spp.

2. RACEME: spikelets have pedicels along the axis, eg. *Digitaria*, *Paspalum* and *Brachiaria* spp. Racemes are more frequent than spikes.

3. PANICLE: spikelets have short stalks on a branched inflorescence with a central axis and a number of side branches. The panicle may be open and loose (*Panicum maximum*), contracted (*Sporobolus* and *Sorghum* spp.) or spike-like and dense (*Cenchrus ciliaris* and *Setaria anceps*) or 'false-spike' when the branches of spike-like panicle are concealed by the spikelets (*Pennisetum purpureum*).



THE FLOWER

The floral organs consists of the gynoecium (female parts), androecium consists of three or one to six, stamens. Each stamen has a slender filament supporting a two celled slender filament supporting a two-celled anther, which consists of the pollen grains. Anther are coloured yellow, purple reddish or may be mottled.

The lodicules base of the flower, outside the stamens. The flowers of most grasses are perfect (hermaphrodite) i.e the florets have both stamens and pistils except members of the tribe maydese. *Zea mays*-male and female separated on the same plant.

ture is such that...





Fig. 3.2 Flowering of the grass plant: (a) Spikelet showing arrangement of florets; (b) Floret opening at blooming time; (c) Typical grass flower showing essential reproductive organs.

Grass plant



BOTANY OF LEGUMES

Legumes are dicotyledonous, i.e. the embryo consists of two cotyledons or seed-leaves). The legume family is sometimes divided into three groups or subfamilies: Mimosoideae, woody plants and herbs with regular flowers, caesalpinoideae, plants with irregular flowers; papilionaceae, herbaceous and woody plants with a distinctive papilionate or butterfly shaped flower. Most of the forage and economically important legumes belongs to papilionaceae family. Legumes may be annuals, biennials or perennials.

A. Vegetative organs

1. Aerial parts

There are distinct morphological differences among the legumes but general characteristics of some plant are similar and rather uniform. The above ground portion consists of a main stem with axillary branches, usually compound leaves, stipules and inflorescences. Tillers sometimes arise from the basal portion of the stem (crown) and stems also develop axillary branches. The stems are jointed, with nodes and internodes, and are usually hollow, except at the nodes. They may be covered with hairs or may be glabrous. Herbaceous stems contains chlorophyll. The leaves contains a common leaf stalk (petiole), with 3 or more leaflets, each with its own stalk

(petiolule). The leaves could be 'palmately' compound i. e leaflets directly attached to the end of the petiole e.g. *Centrosema pubescense* or 'pinnately'

Compound when the petiole extends into a long slender structure with leaflets e.g. *Clitoria ternatea*. Some have leaflets modified to tendrils e.g. *Lathyrus spp.* Presence of *pulvinus* is the characteristic feature of legume family.

Stipules are leaf-like outgrowths found at the base of the main leaf stalk, vary in shape and size and used for identification of species. The leaflets and stipules may be smooth or possess hairs. The veins on the leaves are netted pattern unlike parallel venation of grasses.

2. Subterranean parts

The roots system of most legumes consists principally of an actively growing primary roots and its branches (secondary). The primary roots may penetrate the soil to a depth of 6-8m e. g Lucerne. The roots of many leguminous plants become infected by bacteria of the species *Rhizobium*, Which grows and multiply, forming nodules which differ in size, shape and arrangement on the roots.

B. Reproductive organs

Inflorescence

The Mimosoideae produces flowers in dense heads or small globular, spike – like inflorescences, and commonly has the floral parts arranged in the sets of four, They are rendered conspicuous by the long, coloured filaments of the numerous stamens. e. g *Leucaena leucocephala* and *Acacia spp.*


The caesalpinoideae flowers appears in clusters or racemes, with overlapping petals .The stamens are usually separated e. g *cassia spp.*, *Ceratonia spp.* and *Gleditschia*.

The flowers of papilionaceae are arranged in racemes as in *Desmodium spp.* in heads as in *Trifolium spp.* or spike-like racemes as in *Medicago sativa*. There is a central axis, along with the individual flowers develop. Each flower has its own short stalk or peduncle. The inflorescence may be terminal or auxiliary.


The flower

The corolla consists of five petals of three distinct kinds : 'standards' or 'banner' uppermost or outer petal, largest and most showy; two wing petals, with slender stalks called the claw , and an expanded portion; keel, two petals folded together, partially concealed by the wing petals, the expanded portions more or less united at the outer margin into a boat-shaped structure.

The calyx with five teeth forms a tube at the base of the corolla. The keel encloses the stamens and pistil .The androecium consists of the ten stamens, the filaments of which may be united. The Staminal tube surrounds a superior ovary, an elongated structure comprising one carpel with one ovule or a single row or several ovules. A bent style surmounts the ovary and the stylar tip broadens into the stigmatic surface. The nectar resides at the bottom of the corolla.



The pea flowers are pale yellow with purple or maroon speckles.
Photo by R. P. Ellis courtesy Ecoport
(source URL: <http://www.ecoport.org>)



Habit. A soft wooded tree up to 6 m tall.
Photo by R. P. Ellis courtesy Ecoport

Fruit and Seed

The ripened ovary forms a fruit of variable shape, called legume or pod. Some seeds shatters – dehiscent e. g *Macroptilium atropurpureum* and *Centrosema pubescens* while some are non-dehiscent e. g *Crotalaria* spp. Pods can the reserve food is stored in the cotyledons. Pods can be glabrous or covered with hairs. Each seeds is enclosed in the testa or seedcoat. Hilum marks place of attachment o the ovary walls. The legume seed has no endosperm.

Growth habit

1. Bush – type – a central stalk with side branches appearing along the main stem and with axillary branches developing *Cajanus cajan*; *Desmodium tortuosum*.
2. Bunch – type- a single crown from which several stems and new tillers arise, making it difficult to identify a main stems, stems can be erect or decumbents e. g. *Stylosanthus guianensis* and *Medicago sativa*.
3. Creeping – the stems trail over the ground surface e.g *Calopogonium mucunoides*, *Macroptilium atropurpureum*, some vigna spp.
4. Scrambling – many of the creeping plants climb onto and grow over upright objects. Some are also twining and encircle upright objects e. g *Centrosema pubescens*, *Pueraria phaseoloides*.
5. Rosette - a vegetative form of some perennials developed after flowering or the onset of cool weather e. g *Medicago sativa* and *Trifolium pratense* at the higher elevations of the tropics.





Figure 42 *Leucaena leucocephala* subsp. *glabrata*

Pinnately Compound Leaf	Branch Bearing Simple Leaves
1.It never bears a terminal bud.	It always ends in a terminal bud.
2.It arises from a node.	It develops in the axil of a leaf.
3.Leaflets have no axillary buds.	The leaves borne on a branch have bud in their axil.
4.The rachis of a compound leaf has no nodes and internodes.	The branch bearing simple leaves is always provided with nodes and internodes.
5.Stipules, if present, are seen at the base of the compound leaf, not at base of leaflets.	Stipules, if present, are seen at the base of every leaf.

Pinnately Compound Leaf (ash)



Botanically, the term describes an arrangement of discrete structures (such as leaflets, veins, lobes, branches, or appendages) arising at multiple points along a common axis. For example, once-divided leaf blades having leaflets arranged on both sides of a "[rachis]" are 'pinnately compound' leaves. Plants with pinnate leaves are sometimes colloquially called "feather-leaved".

'paripinnate': pinnately-compound leaves in which leaflets are born in pairs along the rachis without a single terminal leaflet; also called "even-pinnate".

'imparipinnate': pinnately-compound leaves in which there is a lone terminal leaflet rather than a terminal pair of leaflets; also called "odd-pinnate".

'bipinnate': pinnately compound leaves in which the leaflets are themselves pinnately-compound; also called "twice-pinnate".


'tripinnate': pinnately compound leaves in which the leaflets are themselves bipinnate; also called "thrice-pinnate".

'tetrapinnate': pinnately compound leaves in which the leaflets are themselves tripinnate.




Palmately compound leaves

- Based on the type of leaf blade (lamina) leaves are classified into two types namely simple leaves and compound leaves. All leaflets of a compound leaf are borne on common axis and lack axillary buds in their axils.
- In palmately compound leaf all the leaves are attached at a common point i.e. the tip of petiole. This leaves are into three or more lobes (leaflets) and appears like the fingers from the palm of a hand.



Unifoliate palmately compound leaves



Bifoliate palmately compound leaves




Trifoliate palmately compound leaves



Quadrifoliate palmately compound leaves



Multifoliate palmately compound leaves



Palmately Compound Leaf (Horsechestnut)

5 cm

Reproductive organs

Inflorescence

The Mimosoideae produce flowers in dense heads or small globular, spike-like inflorescences, and commonly has the floral parts arranged in the sets of four. They are rendered conspicuous by the long, coloured filaments of the numerous stamens. e.g. *Leucaena leucocephala* and *Acacia spp.*

The caesalpinoideae flowers appear in clusters or racemes, with overlapping petals. The stamens are usually separated e.g. *Cassia spp.*, *Ceratonia spp.* and *Gleditschia*.

The flowers of papilionaceae are arranged in racemes as in *Desmodium spp.* in heads as in *Trifolium spp.* or spike-like racemes as in *Medicago sativa*. There is a central axis, along with the individual flowers develop. Each flower has its own short stalk or peduncle. The inflorescence may be terminal or axillary.

The flower

The corolla consists of five petals of three distinct kinds : 'standards' or 'banner' uppermost or outer petal, largest and most showy; two wing petals, with slender stalks called the claw , and an expanded portion; keel, two petals folded together, partially concealed by the wing petals, the expanded portions more or less united at the outer margin into a boat-shaped structure.

The calyx with five teeth forms a tube at the base of the corolla.

The keel encloses the stamens and pistil. The androecium consists of the ten stamens, the filaments of which may be united.

The Staminal tube surrounds a superior ovary, an elongated structure comprising one carpel with one ovule or a single row or several ovules.

A bent style surmounts the ovary and the stylar tip broadens into the stigmatic surface. The nectar resides at the bottom of the corolla tube.

Fruit and Seed

The ripened ovary forms a fruit of variable shape, called legume or pod. Some seeds shatters - dehiscent e.g *Macroptilium atropurpureum* and *Centrosema pubescens* while some are non-dehiscent e.g *Crotolaria* spp.

Pods can the reserve food is stored in the cotyledons.

be glabrous or covered with hairs. Each seeds is enclosed in the testa or seedcoat. Hilum marks place of attachment to the ovary walls. The legume seed has no endosperm.

Subterranean parts


The roots system of most legumes consists principally of an actively growing primary roots and its branches(secondary). The primary roots may penetrate the soil to a depth of 6-8cm e.g Lucerne. The roots of many leguminous plants become infected by bacteria of the species *Rhizobium*, Which grows and multiply, forming nodules which differ in size, shape and arrangement on the roots.

Growth habit

1. Bush – type – a central stalk with side branches appearing along the main stem and with axillary branches developing *Cajanus cajan*; *Desmodium tortuosum*.
2. Bunch – type- a single crown from which several stems and new tillers arise, making it difficult to identify a main stems, stems can be erect or decumbents e.g *Stylosanthus guianensis* and *Medicago sativa*.
3. Creeping – the stems trail over the ground surface e.g *Calopogonium mucunoides*, *Macroptilium atropurpureum*, some vigna spp.

4. Scrambling – many of the creeping plants climb onto and grow over upright objects. Some are also twining and encircle upright objects e.g *Centrosema pubescens*, *Pueraria phaseoloides*.

5. Rosette- a vegetative form of some perennials developed after flowering or the onset of cool weather e.g *Medicago sativa* and *Trifolium pratense* at the higher elevations of the tropics.



Classification

Grasses

Graminae (Poaceae) is a large botanical family with about 10,000 spp. grouped into some 650 genera and genera into 50-60 tribes; with sub families of 2 to 12.

3 group are of interest- Festucoid group-temperate grasses; Panicoid group- tropical and subtropical and Chloricoid group- few tropical cultivated and a number of valuable wild grasses the tropical and warmer areas of North American .

Examples of festucoid- Tribes of Triticae (Agropyron spp) Festuceae (Festuca, Dactylis, Lolium, poa) Bromeae (Bromus), Aveneae (Avena, Arrhenetherum) Agrostideae (Agrostis, Alopecurus, Phleum)

Panicoid- Paniceae or mellinidae (*Panicum, Brachiaria, Digitaria, Mellinis, Pennisetum, Cenchrus*), Andropogoneae (*Andropogon, Hyparrhenia, Sorghum, Lasiurus, Themedia*) and Maideae (*Zea, Euchlaena, Tripsacum*)

Chloridoid- chlorideae (*Chloris, Cynodon*) and Eragrosteae (*Eragrostis, Dactyloctenium, Eleusine*)

Legume

Order- Leguminosae (Fabaceae) family is divided into three distinct groups or families- Mimosoideae, Caesalpinoideae, Papilionoideae and subfamilies- Mimosaceae, Caesalpinaceae and Papilionaceae of the botanical order Leguminosae.

Only two spp of mimosoideae are important to us – *Leucaena leucocephala* and *Desmanthus virgatus*, although a number of shrubs and trees are browse plant of *Acacia* spp.

A number of Caesalpinioideae are cultivated for fodder and only very few are used as natural browse plants.

Species of Papilionoideae are widely grown as pasture or fodder crops and are of considerable important for natural grazing or browsing.

It is a large subfamily with 200 genera and some 12,000 spp. distributed throughout the world.


Leguminosae in general are of tropical origin with Caesalpinaceae as the most primitive type .

17 tribes of Papilionoideae are recognized .

With important ones as Indigofera, Aeschynomeneae, Sesbanieae, Genisteae, Psoraleae and Trifolieae but majority of legume in cultivation belongs to the tribes – Stylosantheae, Desmodieae and Phaseoleae. The above tribes especially the last three are distributed all over the world, with Desmodieae and Phaseoleae concentrated in Latin American.

ESSENTIAL SOIL NUTRIENTS FOR PASTURE PRODUCTION



- **NITROGEN:** Legumes may supply substantial quantities of N to the soil through fixation by symbiotic relationship with rhizobia.
- Legumes can absorb nitrogen from that which is available in the soil.
- Nitrogen made available to plants from the soil is derived from applied fertilizer, precipitation and non-symbiotic fixation.
- Negative effects of N fertilizer in grass-legume sward






Nitrogen Sources	
Source Material	% Nitrogen Content
Anhydrous ammonia	82%
Aqua ammonia	20% to 25%
Ammonium nitrate	33.5% to 34%
Ammonium nitrate-sulfate	26%
Ammonium nitrate/lime	20.5%
Ammonium sulfate	21%
Urea-ammonium nitrate solution	28% to 32%
Ammonium chloride	26%
Urea	46%
Monoammonium phosphate	10% to 11%
Diammonium phosphate	18%
Sodium nitrate	16%
Potassium nitrate	13%
Calcium nitrate	15.5%
Sulfur-coated urea	39%
Urea-formaldehyde	38%

Lecture 1: Fertilizers and Soil Fertility





- Nitrogen is a mobile nutrient.
- PHOSPHORUS: The P requirement of grasses unlike that of N depends more on soil properties than on the grass species.
- P is likely to be limiting factor when grasses are established on land not previously cultivated and fertilized with phosphate.
- P is less mobile in soils than N and K and remains fairly close to the area in which it is placed.
- Molybdenum: It is essential in the formation of nitrate reductase and plays an important role in the symbiotic fixation of nitrogen
- Molybdenum deficiency resembles N starvation and plant symptoms are likely to occur when legumes grow on soils low in available N.
- Potassium: The influence of K on productivity of pasture species or cultivars varies with the requirement of this nutrient for plant growth, ability of plants to extract the nutrient from the soil.
- K Deficiency: a K content of less than 1.0-1.5% in 60 day old grass in an intensive management system will show k deficiency.
- The response of tropical legumes to applied k fertilizers has been variable.
- Sulphur: S deficient soils generally exhibit high allophane or oxides, low in organic matter and often sandy soils may be deficient in S where annual burning occurs.

- Boron: Deficiency not common but was corrected with 25-30kg/ha of borax when noted in lucerne growing on certain soils
- Aluminium: Al toxicity is likely to occur in tropical soils with a pH of 4.5 or less. The effect on plant is largely through root damage with restriction of terminal elongation and branching along with diminished nodulation.
- Manganese toxicity is sometimes compounded with Al toxicity since both ions are progressively released with increasing soil acidity.
- Calcium: Limestone (CaCO_3) application to pastures in the tropics should be considered as a source of nutrient Ca rather than a soil amendment.
- If lime application is needed for maximum productivity of legumes
- Liming decreased the manganese content of soil and herbage but did not alter P or Mg contents.
- Ca can be transported from lower soil depths to upper layers by roots